

**Method for producing an active heterodimeric AMV-RT in  
prokaryotic cells**

---

The invention concerns a method for producing a recombinant active heterodimeric AMV-RT by expressing one or several DNA sequences coding for the  $\alpha$ - and/or  $\beta$ -subunit(s) of the AMV-RT in prokaryotic cells under certain growth and induction conditions.

The discovery of the reverse transcriptases in the seventies disproved the "central dogma" of molecular biology on the information transfer from DNA via RNA to protein as a unidirectional process (Termin H. and Mizutani S., 1970 Nature 226:1211-1213; Baltimore D., 1970, Nature 226:1209-1211). The enzymatic characterization of these RNA-dependent DNA polymerases is the basis for current understanding on the amplification cycle of RNA viruses and thus also on the development and spread of diseases that are caused by this type of virus (cancer, AIDS etc.).

However, reverse transcriptases are also a tool for molecular biologists for the synthesis, amplification and cloning of cDNAs (RT-PCR). This technology allows a simplified and accelerated examination of gene expression in eukaryotic cells. After isolating the total mRNA from cell extracts or tissues, the mRNA is translated back into cDNA by the reverse transcriptase and amplified by the subsequent PCR step to enable cloning and characterization. Consequently it is not necessary to, on the one hand, elucidate the intron and

exon structures of the genes but, on the other hand, it is also possible to examine gene expression in the cell during various life cycles or during the development of diseases (such as cancer).

Reverse transcriptases (RT) from three different retroviruses have hitherto been closely examined: The RT from *Moloney Murine Leukemia Virus* (M-MLV). This enzyme consists of a single subunit with a molecular weight of 78 kDa (Prasad V.R., 1993 reviewed in *Reverse Transcriptase*, Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press, 135). In addition an RT from *Human Immunodeficiency Virus* (HIV) is known. This RT is a heterodimer that is composed of two subunits p66 and p51, the p51 subunit being formed by proteolytic cleavage of p66 (Le Grice S.F.J., 1993 reviewed in *Reverse Transcriptase*, Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press, 163). In addition RTs are known from *Avian Sarcoma-Leukosis Virus* (ASLV). The RT obtainable from *Avian Myeloblastosis Virus* (AMV) also belongs to the ASLV family. This RT is also a heterodimer that is composed of an  $\alpha$ -chain with a molecular weight of ca. 63 kDa and a  $\beta$ -chain with a molecular weight of ca. 95 kDa. In this case the  $\alpha$ -chain is also formed by proteolytic processing of the  $\beta$ -chain (Golomb M. and Grandgenett D., 1979, *J. Biol. Chem.* **254**: 1606-1613; Weiss R. et al., eds. 1984, *Molecular Biology of tumor viruses*, 2<sup>nd</sup> edition: RNA tumor viruses 1/text. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York).

Whereas the M-MLV-RT is expressed in *E. coli* as a monomer and the HIV-RT as a heterodimer, it has so far not been possible to express the AMV-RT as an active or soluble heterodimer in *E. coli* or other prokaryotes to a

satisfactory degree. Although according to WO 00/42199 certain RT variants are expressed in *E. coli* or preferably in eukaryotic insect cells, the desired RT that is obtained in this process mainly consists (ca. 90 %) of an insoluble component.

In addition it is difficult to measure a recombinant AMV-RT in crude cell extracts of *E. coli* since, on the one hand, RNA templates are degraded by intrinsic *E. coli* RNases and, on the other hand, *E. coli* strains have a DNA polymerase which also has an RT activity in addition to the DNA polymerase activity (Ricchetti, M. and Huc, H., 1993, *EMBO J.* 12 (2), 387-396). Hence this intrinsic *E. coli* RT activity considerably interferes with the determination of the activity of the recombinant AMV-RT in crude *E. coli* extracts and in fractions from the purification.

Hence the object of the present invention is to provide a recombinant active heterodimeric AMV-RT in adequate amounts.

The object is achieved by a method for producing an active heterodimeric AMV-RT in prokaryotic host cells wherein one or several DNA sequence(s) which code for the  $\alpha$  and  $\beta$  subunit or chain of the AMV-RT, are cloned into expression plasmids, the expression plasmids are transformed in prokaryotic cells, the expression of the heterodimeric AMV-RT is induced and the recombinant heterodimeric AMV-RT is purified i.e. isolated from the cells. Suitable genes and DNA sequences are, among others, those which only code for one of the AMV-RT subunits. A portion or the expression product can subsequently be converted by certain measures, such as

proteolytic cleavage of the  $\beta$ -chain, into the other subunit. The sequences SEQ ID NO:4 and SEQ ID NO:5 have proven to be particularly suitable for the method according to the invention which generates an active heterodimeric AMV-RT composed of the subunits SEQ ID NO:6 and SEQ ID NO:7.

The structural genes and DNA sequences coding for the subunits of the AMV-RT can either be cloned on different, separate expression plasmids or on one expression plasmid, optionally in the presence of so-called helper plasmids, and expressed in a suitable host cell. Suitable expression plasmids are for example pDS, pKK177-3 or pKKT5. The plasmid pKKT5 in which the respective structural genes are inserted under the control of the T5 promoter is preferred according to the invention. Other potential promoters, which are preferably IPTG-inducible promoters, are for example the lac, lac UV5 or tac promoter. Alternative helper plasmids such as the plasmid pUBS520 and suitable selection markers such as ampicillin or kanamycin are in principle known to a person skilled in the art.

The expression plasmids and optionally other helper plasmids are transformed into a suitable prokaryotic host cell. According to the invention it is preferable to use an *E. coli* strain such as *E. coli* K12 C600, DH5 $\alpha$ , LE392, JM83, JM105, NM522, M15, RR1 $\Delta$ 15, UT5600, TG1, A1200 or the strains *E. coli* B, BL21, HB101. The *E. coli* strain LE392 is particularly preferred according to the invention.

The expression of the heterodimeric AMV-RT can be induced by various measures. In particular certain

growth and induction conditions have positive effects on the expression of active AMV-RT. A growth temperature in the range of 10° to 25°C combined with a low inducer concentration has proven to be advantageous according to the invention. A growth temperature of about 15°C and an inducer concentration between 0.1 and 0.5 mM, preferably of about 0.15 mM, have proven to be particularly suitable. IPTG (isopropyl- $\beta$ -D-thiogalactopyranoside) or lactose are preferably used according to the invention as the inducer.

Furthermore it turned out that the soluble expression of AMV-RT in prokaryotic cells can be increased by the co-expression of helper genes. Potential helper genes are in particular the trpT gene which codes for the tryptophan tRNA. In addition chaperone genes are suitable for soluble expression such as the genes coding for GroEL and GroES, GrpE, ClpB, Dnak and DnaJ. The genes for one or several chaperones are then preferably located on a helper plasmid with an inducible promoter; the genes which code for the chaperones GroEL and GroES are under the control of a constitutive promoter on the expression plasmid on which the structural genes for the  $\alpha$  and/or  $\beta$  chain are also located. However, it is particularly preferred according to the invention when the genes coding for GroEL and GroES are cloned on the expression plasmid which carries the genes for the  $\alpha$ - and  $\beta$ -chain and the genes coding for Dnak, DnaJ, GrpE and ClpB are cloned on a helper plasmid.

In addition to methods that are generally known to a person skilled in the art, it is especially advantageous to use affinity chromatography materials such as metal ion chelating materials or cation exchangers to purify and isolate the recombinant heterodimeric AMV-RT from

the cell extract. It is particularly advantageous for the purification of the AMV-RT for the expression products, i.e. the  $\alpha$ - as well as the  $\beta$ -chain to be fused with peptide sequences that are able to reversibly bind to particular column materials such as cation exchangers, metal ion chelating materials such as nickel, copper or zinc nitriloacetic acid (NTA) resins. Peptide sequences that are suitable according to the invention can have from two to about 100 amino acids or amino acid derivatives. Peptide sequences which are composed of two to ten amino acids, e.g. arginine residues or histidine residues, have proven to be particularly suitable for the invention. In addition it has also proven to be particularly advantageous to use such peptide sequences comprising eight arginine or six histidine residues. In addition commercially available peptide sequences such as Strep-tag<sup>®</sup> (IBA GmbH, Göttingen/Germany) or GST-tag (Pharmacia, Uppsala/Sweden) are also suitable for the method according to the invention.

The invention is further elucidated by the following examples:

**1. Example:**

**Isolation of genes which code for the  $\alpha$ - chain and  $\beta$ -chain**

The data bank sequence (MEDLINE ID 94366722, Baluda et al., 1994) was used to design oligonucleotide primers for the isolation of the  $\beta$ -chain (see SEQ ID NO:1 and 2). A *EcoRI* restriction endonuclease cleavage site was incorporated at the 5' end and a *PstI* restriction cleavage site was incorporated at the 3' end for the

subsequent cloning into vectors. In addition a further 3' primer was designed (see SEQ ID NO:3) which enables the isolation of the  $\alpha$ -chain. Both chains were fished by means of PCR from a virus lysate (ATCC VR-265) by means of RT-PCR as well as from an *E. coli* clone (ATCC 31990) which carries the  $\beta$ -chain on a plasmid. The PCR mixtures were applied to a 1 % agarose gel, the PCR fragments of ca. 1715 bp for the  $\alpha$ -chain and ca. 2570 bp for the  $\beta$ -chain were isolated from the agarose gel (QIAEX II, Gel Extraction Kit, Qiagen/Germany), cleaved with the restriction endonucleases mentioned above and cloned into a vector fragment of pUC19 that had also been linearized with *EcoRI* and *PstI* and isolated. For this 1  $\mu$ l (20 ng) vector fragment and 3  $\mu$ l (100 ng) PCR fragment, 1  $\mu$ l 10 x ligase buffer (Maniatis et al., 1989 Molecular Cloning: A Laboratory Manual, second Edition, Cold Spring Harbor Laboratory Press NY (USA), vol. 27), 1  $\mu$ l T4 DNA ligase, 4  $\mu$ l sterile H<sub>2</sub>O<sub>bidistilled</sub> were pipetted, carefully mixed and incubated overnight at 16°C. The cloned genes were subsequently examined by means of restriction analysis and sequencing. The sequences are shown in SEQ ID NO:4 ( $\alpha$ -chain) and SEQ ID NO: 5 ( $\beta$ -chain).

Comparison with the data bank sequence (MEDLINE ID 94366722, Baluda, M.A., and Reddy, E.P., 1994, Oncogene 9:2761-2774) yielded a homology of 98.8 % at the DNA level for the  $\alpha$ -chain as well as for the  $\beta$ -chain. When the resulting amino acid sequences are compared, it becomes apparent that most substitutions at the DNA level are so-called silent mutations i.e. do not lead to amino acid substitutions. Only three base substitutions also resulted in amino acid substitutions but they are found reproducibly in each isolated PCR product. These are the substitutions Arg273Met, Arg304Gln and

Asp495Glu. The amino acid sequences of both chains are shown in SEQ ID NO:6 ( $\alpha$ -chain) and SEQ ID NO:7 ( $\beta$ -chain).

## 2. Example:

### Expression of the $\alpha$ -chain and $\beta$ -chain without fused peptide sequences (tags)

#### 2.1. Construction of the expression plasmids pAMV- $\alpha$ and pAMV- $\beta$

In order to express the AMV-RT, the genes for both chains were cloned separately into expression vectors in such a manner that the structural genes were each inserted in the correct orientation under the control of the T5 promoter. For this the respective structural gene for the  $\alpha$ -chain and the  $\beta$ -chain were cut out of the plasmid pUC19 by *EcoRI* and *PstI*, the restriction mixtures were separated by agarose gel electrophoresis and the 1715 bp fragment of the  $\alpha$ -chain and the 2570 bp fragment of the  $\beta$ -chain were isolated from the agarose gel. The expression plasmid pKKT5, which is formed from pKK177-3 (Kopetzki et al., 1989, *Mol. Gen. Genet.* **216**: 149-155) by replacing the *tac* promoter with the T5 promoter from pDS (Bujard et al., 1987, *Methods Enzymol.* **155**: 416-433), was used for the expression. The *EcoRI* restriction endonuclease cleavage site in the sequence of the T5 promoter was removed by two point mutations. The resulting expression plasmid was cut with *EcoRI* and *PstI* for the insertion of the genes for the AMV-RT, the restriction mixture was separated by agarose gel electrophoresis and the resulting vector fragment of ca. 2500 bp was isolated from the agarose gel. The vector fragment obtained in this manner was separately ligated



as described above with the genes for the  $\alpha$ -chain and the  $\beta$ -chain described in example 1. The correct insertion of the genes was checked by restriction control and sequencing. The resulting plasmid pAMV- $\alpha$  and pAMV- $\beta$  was firstly separately cotransformed with the helper plasmid pUBS520 for expression control in various *E. coli* strains. It is conceivable in this case that the  $\alpha$ -chain and the  $\beta$ -chain could be separately expressed in order to obtain  $\alpha\alpha$ - and  $\beta\beta$ -homodimers. The helper plasmid pUBS520 (Brinkmann et al., 1989, Gene 85: 109-114) carries inter alia the *lacI<sup>q</sup>* gene which codes for the lac repressor and the *dnaY* gene which codes for the rare tRNA<sup>Arg</sup> in *E. coli* which recognizes the codons AGA and AGG (Garcia et al., 1986, Cell 45: 453-459). The kanamycin resistance gene from the transposon TN903 was used as a selection marker.

## 2.2 Separate transformation of the expression plasmids pAMV- $\alpha$ and pAMV- $\beta$ in *E. coli*

Competent cells of various *E. coli* strains were prepared according to the method of Hanahan (*J. Mol. Biol.* 1983, vol. 166, 557). 200  $\mu$ l of *E. coli* LE392 cells prepared in this manner were admixed with 20 ng isolated expression plasmid pAMV- $\alpha$  DNA or pAMV- $\beta$  DNA and 40 ng helper plasmid DNA. After 30 min incubation on ice a heat shock (90 sec. at 42°C) was carried out. Subsequently the cells were transferred to 1 ml LB medium and incubated for 1 hour at 37°C in LB medium for the phenotypic expression. Aliquots of this transformation mixture were plated out on LB plates containing ampicillin and kanamycin as selection markers and incubated for 15 hours at 37°C.

### 2.3 Expression of the gene for the $\alpha$ -chain in *E. coli*

In order to express the gene which codes for the  $\alpha$ -chain of the AMV-RT, plasmid-containing clones were inoculated in 3 ml LB<sub>ampkan</sub> medium and incubated at 30°C in a shaker. At an optical density of 0.5 (measured at 550 nm, OD<sub>550</sub>) the cells were induced with 0.5 mM IPTG and incubated for 4 h at 30°C in a shaker. Subsequently the optical density of the individual expression clones was determined, an aliquot which corresponded to an OD<sub>550</sub> of 5.0/ml was removed and the cells were centrifuged (10 min, 6000 rpm, 4°C). The cell pellet was resuspended in 400  $\mu$ l TE buffer (50 mM TRIS/50 mM EDTA, pH 8.0), the cells were disrupted by ultrasound and the soluble protein fraction was separated from the insoluble protein fraction by centrifugation (10 min, 14000 rpm, 4°C). Application buffer containing SDS and  $\beta$ -mercaptoethanol was added to all fractions and the proteins were denatured by boiling (5 min 100°C). Afterwards 10  $\mu$ l of each was analysed by means of an analytical SDS gel (10 %) (Laemmli U.K., 1970, *Nature* 227: 555-557).

Analysis of the SDS gel shows a clear overexpression of the  $\alpha$ -chain. A strongly overexpressed additional band is seen at ca. 63 kDa which is not observed with the non-induced control clones or the induced control clones which do not contain plasmid. A small portion of the overexpressed  $\alpha$ -chain appears in the soluble protein fraction whereas the major amount is formed as an insoluble expressed protein.

### 2.4 Expression of the gene of the $\beta$ -chain in *E. coli*

In order to express the gene which codes for the  $\beta$ -chain

of the AMV-RT, 3 ml LB<sub>ampkan</sub> medium was inoculated with plasmid-containing clones and incubated at 30°C in a shaker. At an OD<sub>550nm</sub> of 0.5 the cells were induced with 0.5 mM IPTG and incubated for 4 h at 30°C in a shaker. Subsequently the optical density of the individual expression clones was determined, an aliquot which corresponded to an OD<sub>550</sub> of 5.0/ml was removed and the cells were centrifuged (10 min, 6000 rpm, 4°C). The cell pellet was resuspended in 400 µl TE buffer (50 mM TRIS/50 mM EDTA, pH 8.0), the cells were disrupted by ultrasound and the soluble protein fraction was separated from the insoluble protein fraction by centrifugation (10 min, 14000 rpm, 4°C). Application buffer containing SDS and β-mercaptoethanol was added to all fractions and the proteins were denatured by boiling (5 min 100°C). Afterwards 10 µl of each was analysed by means of an analytical SDS gel (8 %) (Laemmli U.K., 1970, *Nature* **227**: 555-557).

Analysis of the SDS gel shows a clear overexpression of the β-chain. A strongly overexpressed additional band is seen at ca. 95 kDa which is not observed with the non-induced control clones or the induced control clones which do not contain plasmid. The majority of the overexpressed β-chain appears in the insoluble protein fraction, however, a slight overexpression is also seen in the soluble protein fraction.

## **2.5 Expression of both chains on separate plasmids in a cell**

In order to express both chains in one cell, the *lacI<sup>q</sup>* expression cassette and the *dnaY* expression cassette must firstly be recloned from the helper plasmid pUBS520

onto the expression plasmids. The *lacI<sup>q</sup>* expression cassette was cloned onto pAMV- $\alpha$  and the *dnaY* expression cassette was cloned onto the expression plasmid pAMV- $\beta$ . In order to ensure a stable multiplication of the expression plasmids, the ampicillin resistance gene from pAMV- $\alpha$  was replaced by the kanamycin resistance gene from pUBS520. The resulting expression plasmids pAMV- $\alpha$  *lacI<sup>q</sup>* and pAMV- $\beta$  *dnaY* were subsequently cotransformed in various *E. coli* expression strains.

In order to express the genes which code for the  $\alpha$ -chain and the  $\beta$ -chain of the AMV-RT, 3 ml LB<sub>ampkan</sub> medium was inoculated with plasmid-containing clones and incubated at 30°C in a shaker. At an OD<sub>550nm</sub> of 0.5 the cells were induced with 0.5 mM IPTG and incubated for 4 h at 30°C in a shaker. Subsequently the optical density of the individual expression clones was determined, an aliquot which corresponded to an OD<sub>550</sub> of 5.0/ml was removed and the cells were centrifuged (10 min, 6000 rpm, 4°C). The cell pellet was resuspended in 400  $\mu$ l TE buffer (50 mM TRIS/50 mM EDTA, pH 8.0), the cells were disrupted by ultrasound and the soluble protein fraction was separated from the insoluble protein fraction by centrifugation (10 min, 14000 rpm, 4°C). Application buffer containing SDS and  $\beta$ -mercaptoethanol was added to all fractions and the proteins were denatured by boiling (5 min 100°C). Afterwards 10  $\mu$ l of each was analysed by means of an analytical SDS gel (8 %) (Laemmli U.K., 1970, *Nature* **227**: 555-557).

Analysis of the SDS gel surprisingly shows a clear overexpression of the  $\alpha$ - and  $\beta$ -chain. Strongly overexpressed additional bands are seen at ca. 63 kDa and ca. 95 kDa which are not observed with the non-induced control clones or the induced control clones which do not

contain plasmid. The distribution of the bands in the soluble and insoluble fraction is like that of the experiments in which both chains were expressed separately. The expression output of both chains is overall somewhat less than for separate expression.

### 3. Example:

**Expression of the  $\alpha$ -chain and  $\beta$ -chain with fused tags to simplify the purification**

#### 3.1 Production of various fusion proteins

In order to efficiently purify the recombinant AMV-RT heterodimers, suitable peptide sequences, so-called tags were fused to the 5' end of both chains. Tags enable affinity chromatographies to be carried out. A series of two affinity chromatographies which are each specific for one of the two tags additionally allows the isolation of pure heterodimers (Wende W. et al., 1996, *Biol. Chem.* **377**, 625-632). Appropriate primer designs were used to attach eight arginine residues to the  $\alpha$ -chain and six histidine residues to the  $\beta$ -chain by means of PCR reactions. The sequences of the sense primers are shown in SEQ ID NO:8 (5' primer for the  $\alpha$ -chain) and SEQ ID NO:9 (5' primer for the  $\beta$ -chain). The oligonucleotides of SEQ ID NO:2 ( $\beta$ -chain) and SEQ ID NO:3 ( $\alpha$ -chain) which had already been used for gene isolation were used as antisense primers.

The PCR mixtures were applied to a 1 % agarose gel, the PCR fragments of 1739 bp for the  $\alpha$ -chain and 2597 bp for the  $\beta$ -chain were isolated from the agarose gel (QIAEX II, Gel Extraction Kit, Qiagen, Germany), cleaved with the restriction endonucleases *EcoRI* and *PstI* and cloned

into a vector fragment of the preferred expression plasmid that had also been linearized with *EcoRI* and *PstI* and isolated. For this 1  $\mu$ l (20 ng) vector fragment and 3  $\mu$ l (100 ng) PCR fragment, 1  $\mu$ l 10 x ligase buffer (Maniatis *et al.*, 1989 Molecular Cloning: A Laboratory Manual, second Edition, Cold Spring Harbor Laboratory Press NY (USA), vol. 27), 1  $\mu$ l T4 DNA ligase, 4  $\mu$ l sterile H<sub>2</sub>O<sub>bidistilled</sub> were pipetted, carefully mixed and incubated overnight at 16°C. The cloned genes were subsequently examined by means of restriction analysis and sequencing. The resulting expression plasmids were named pAMV- $\alpha_{lacIq}$ -Arg and pAMV- $\beta_{dnaY}$ -His.

### **3.2 Transformation of the expression plasmids pAMV- $\alpha_{lacIq}$ -Arg and pAMV- $\beta_{dnaY}$ -His in various *E. coli* expression strains**

Competent cells of various *E. coli* strains were prepared according to the method of Hanahan (*J. Mol. Biol.* 1983, vol. 166 pp. 557) (see example 2.2).

### **3.3 Expression of both chains with fused tags on separate plasmids in a cell**

In order to express both chains with tags in a cell, various *E. coli* expression strains were cotransformed with the expression plasmids pAMV- $\alpha_{lacIq}$ -Arg and pAMV- $\beta_{dnaY}$ -His.

In order to express the genes which code for the  $\alpha$ -chain with an Arg-tag and the  $\beta$ -chain with an His-tag of the AMV-RT, 3 ml LB<sub>ampkan</sub> medium was inoculated with plasmid-containing clones and incubated at 30°C in a shaker. At an OD<sub>550</sub> of 0.5 the cells were induced with 0.5 mM IPTG and incubated for 4 h at 30°C in a shaker. Subsequently

the optical density of the individual expression clones was determined, an aliquot which corresponded to an OD<sub>550</sub> of 5/ml was removed and the cells were centrifuged (10 min, 6000 rpm, 4°C). The cell pellet was resuspended in 400 µl TE buffer (50 mM TRIS/50 mM EDTA, pH 8.0), the cells were disrupted by ultrasound and the soluble protein fraction was separated from the insoluble protein fraction by centrifugation (10 min, 14000 rpm, 4°C). Application buffer containing SDS and β-mercaptoethanol was added to all fractions and the proteins were denatured by boiling (5 min 100°C). Afterwards 10 µl of each was analysed by means of an 8 % analytical SDS gel (Laemmli U.K., 1970, *Nature* **227**: 555-557).

Analysis of the SDS gel surprisingly shows a clear overexpression of the α- and β-chain. Strongly overexpressed additional bands are seen at ca. 63 kDa and ca. 95 kDa which are not observed with the non-induced control clones or the induced control clones which do not contain plasmid. The distribution of the bands in the soluble and insoluble fraction is like that of the experiments in which both chains were expressed separately without tags in one cell.

### **3.4 Expression of both chains with fused tags on a plasmid**

If the genes for the α- and β-chain of the AMV-RT are distributed on two plasmids, differences in the stability of these plasmids could lead to the production of different amounts of the respective chains and thus to a lower yield of αβ-chain heterodimer. Hence with the exception of the gene for β lactamase, the entire genetic information of the two plasmids pAMV-α<sub>lacIq</sub>-Arg and pAMV-β<sub>dnal-His</sub> was combined on a single plasmid pAMV-

$\alpha\beta$ -1. This plasmid was constructed by inserting the *SspI*-*AflIII* fragment of pAMV- $\beta_{dnaY-His}$  containing the sequence for the T5 promoter, the gene coding for the  $\beta$ -chain with an N-terminal His tag, the sequence for the *rrnB* terminator and the *dnaY* gene, into the *SalI* cleavage site of pAMV- $\alpha_{lacIq-Arg}$  which contains the sequence for the T5 promoter, the gene coding for the  $\alpha$ -chain with an N-terminal Arg-tag, the sequence for the *rrnB* terminator, the kanamycin resistance gene and the *lacIq* gene. For this purpose 1  $\mu$ g each of the expression plasmids pAMV- $\alpha_{lacIq-Arg}$  and pAMV- $\beta_{dnaY-His}$  were cleaved with the restriction endonucleases described above according to the manufacturer's instructions, the restriction mixtures were separated in a 1 % agarose gel and the 4124 bp *SspI*-*AflIII* fragment of pAMV- $\beta_{dnaY-His}$  and the 6024 bp fragment of pAMV- $\alpha_{lacIq-Arg}$  were isolated from the agarose gel (QIAEX II, Gel Extraction Kit, Qiagen/Germany). The non-compatible ends were prepared with Klenow polymerase (Roche Diagnostics GmbH) according to the manufacturer's instructions and the two fragments were ligated together as described above. The resulting new expression plasmid pAMV $\alpha\beta$ -1 was examined by means of restriction analysis.

The correct expression plasmid according to restriction analysis was transformed in the *E. coli* K-12 strain LE392 as described above and subjected to an expression control. The protein content of the cells after 4 hours growth under induced conditions was subsequently examined by means of SDS-PAGE. According to SDS-PAGE analysis the level of the expression output and the relative proportion of soluble and insoluble fractions are comparable to the expression of the genes for the  $\alpha$ - and  $\beta$ -chain on separate plasmids, but the amount of expressed  $\alpha$ - and  $\beta$ -chain appears to be more homogeneous.



Furthermore the Arg-tag of the  $\alpha$ -chain was replaced by a His-tag like that of the  $\beta$ -chain for the purification procedure. For this purpose an intermediate construct pAMV- $\alpha_{lacIq-His}$  was prepared in which the *EcoRI-NheI* fragment from pAMV- $\alpha_{lacIq-Arg}$  was replaced by the *EcoRI-NheI* fragment from pAMV- $\beta_{dnaY-His}$ . Subsequently, like the construction of pAMV $\alpha\beta$ -1, the entire genetic information of the two plasmids pAMV- $\alpha_{lacIq-His}$  and pAMV- $\beta_{dnaY-His}$  with the exception of the gene for  $\beta$  lactamase was combined on a single plasmid pAMV $\alpha\beta$ -2. The new expression vector was named pAMV $\alpha\beta$ -2. Cells were transformed as described above with pAMV $\alpha\beta$ -2 and subjected to an expression control under standard conditions. The expression output was not increased under these conditions.

#### **4. Example:**

##### **Expression optimization**

##### **4.1 Increase of the expression of active AMV-RT by changing the expression conditions**

Particular growth and induction conditions have positive effects on the expression of active AMV-RT. Afterwards the growth temperature was lowered from 30°C to 15°C during the induction phase, the IPTG concentration was reduced from 0.5 mM to 0.15 mM to induce expression and the induction time was increased from 4 h to 26 h. The protein content of the cells after the induction phase was examined as described above by SDS polyacrylamide gel electrophoresis.

Afterwards the total expression yield of  $\alpha$ - and  $\beta$ -chain was, as expected, substantially reduced in the SDS-PAGE analysis, but the content of soluble expressed  $\alpha$ - and  $\beta$ -

chain was considerably increased in comparison to the expression experiments under standard growth and induction conditions. This increase in the expression of active AMV-RT was also confirmed in the subsequent purification and activity determination.

#### 4.2 Increasing the expression of active AMV-RT by coexpression of helper genes

##### 4.2.1. Coexpression of the gene for the tryptophan-tRNA (tRNA<sup>trp</sup>)

One property of the AMV-RT is to use an endogenous cell tRNA for tryptophan (tRNA<sup>trp</sup>) as a primer for the polymerase reaction after infection of a eukaryotic host cell (Leis et al., 1993, in: Reverse Transcriptase, Cold Spring Harbor Monograph Series, eds.: Skala, A.M. and Goff, S.P., Cold Spring Harbor NY (USA), 33-48). However, whether the endogenous *E. coli* tRNA<sup>trp</sup> can be used by the AMV-RT as a primer has not been proven. In *E. coli* the tRNA<sup>trp</sup> is only coded by a single gene *trpT*, the expression of which is adapted to the normal requirements of the cell. In order to exclude a potential deficiency of tRNA<sup>trp</sup> in the cell, the *trpT* gene according to SEQ ID NO:10 was isolated by means of PCR from *E. coli* LE392 (the primers used for this are shown in SEQ ID NO:11 and 12), recleaved with *Pst*I for insertion into pAMV- $\alpha_{lacIq-His}$  and ligated into the vector fragment of pAMV- $\alpha_{lacIq-His}$  that was also linearized with *Pst*I as described above. Clones which have integrated the *trpT* gene at the *Pst*I restriction endonuclease cleavage site were checked by means of restriction analysis and sequencing. In this intermediate construct pAMV- $\alpha_{lacIq-His-trpT}$  the gene for

the  $\alpha$ -chain and the gene for the *E. coli* tRNA<sup>trp</sup> form one transcription unit, the expression of which is regulated by the IPTG-inducible T5 promoter. Subsequently, similarly to the construction of pAMV $\alpha\beta$ -1 or pAMV $\alpha\beta$ -2, the entire genetic information of the two plasmids pAMV- $\alpha_{lacIq}$ -His-*trpT* and pAMV- $\beta_{dnaY}$ -His with the exception of the gene for  $\beta$ -lactamase was combined on a single plasmid pAMV $\alpha\beta$ -3. Cells were transformed as described above with pAMV $\alpha\beta$ -3 and subjected to an expression control using the modified expression conditions. Afterwards the yield of active AMV-RT is significantly increased.

#### 4.2.2. Coexpression of chaperone genes

In *E. coli* there are two main chaperone systems comprising the GroESL machinery and a 4 component system consisting of DnaK, DnaJ, GrpE and ClpB (Kedzierska, 1999). Both systems play an important role in the correct folding of newly formed proteins as well as in the renaturing of proteins that have aggregated as a result of stress (Hartl F.U., 1996, Nature 381, 571-580; Bukau H. and Horwich A.L., 1998, Cell 92, 351-366; Mogk A. et al., EMBO J. 18, 6934-6949; Zolkiewski M., 1999, J. Biol. Chem. 274, 28083-28086; Goloubinoff P. et al., 1999, Proc. Natl. Acad. Sci. USA 96, 13732-13737).

In a first step the *groESL* operon from *E. coli* should be overexpressed in the AMV-RT production strains. For this the *EcoRI-HindIII* fragment from pOF39 (Fayet O., Louarn J.-M., Georgopoulos C., 1986, Mol. Gen. Genet. vol. 202, pp. 335-345) was integrated in the *SspI* cleavage site of the plasmid pAMV- $\beta_{dnaY}$ -His. Non-compatible ends were prepared with Klenow polymerase (Roche Diagnostics)

according to the manufacturer's instructions before ligation. The sequence of *groESL* is shown in SEQ ID NO:13. In this new construct pAMV- $\beta_{dnaY}$ -His-*groESL*, the *groESL* operon forms an artificial transcription unit containing the 3'-situated gene for  $\beta$  lactamase. The expression is then either under the control of the endogenous *bla* constitutive promoter that is now on the 5' side of the *groESL* operon and/or under the control of the  $\sigma^{32}$ -dependent promoter of the *groESL* operon. Subsequently the entire genetic information of the two expression plasmids pAMV- $\alpha_{lacIq}$ -His-*trpT* and pAMV- $\beta_{dnaY}$ -His-*groESL* with the exception of the gene for  $\beta$ -lactamase was again combined as described above on a single plasmid pAMV $\alpha\beta$ -4.

Cells were transformed with pAMV $\alpha\beta$ -4 as described above and subjected to an expression control under the modified expression conditions. The co-overproduction of GroESL results in an increase of the biomass and of the amount of active AMV-RT. Three to four-fold higher values were obtained compared to the previously best production strains after purification and activity testing.

After the co-overproduction of GroESL in the AMV-RT production strains had proven to be a positive measure, the other main chaperone system of *E. coli* was additionally co-overproduced in a second step. In addition to the supposed general advantages of this co-overproduction this could compensate for a disadvantage of the GroESL machinery i.e. its exclusion volume of circa 65 kDa (Deuerling E. et al., 1999, Nature 400, 693-696). This should be particularly important for the correct folding of the  $\beta$ -chain of the AMV-RT (93 kDa) provided it cannot be divided into single domains that

are formed independently of one another. The genes DnaK, DnaJ and GrpE were combined in an artificial operon corresponding to the physiological combination (Diamant S. and Goloubinoff P., 1998, Biochemistry 37, 9688-9694; Pierpaoli E.V. et al., 1998, J. Biol. Chem. 273, 6643-6649), whereas the gene for ClpB forms its own transcription unit. Both transcription units were placed under the control of IPTG-inducible T5 promoters in order to coordinate the expression with the genes for the subunits of the AMV-RT.

For technical reasons the cloning process required a number of intermediate steps on the path to the final construct pCHAP-5. Thus the pKKT5 derivatives pCHAP-1 and pCHAP-2 were firstly constructed. pCHAP-1 contains the genetic information for the *dnaKJ* operon from *E. coli* starting with the start codon for *dnaK* up to the stop codon for *dnaJ*; pCHAP-2 carries the artificial transcription unit from the coding regions of the genes for GrpE and ClpB as an insert; the corresponding DNA fragments were amplified by PCR from the genomic DNA of *E. coli* K12KE392. The sequence of the *dnaKJ* operon is shown in SEQ ID NO:14, the corresponding primers used to isolate the *dnaKJ* operon are shown in SEQ ID NO:15 and 16. The sequence of the *grpE* gene is shown in SEQ ID NO:17, the corresponding primers for the isolation of the *grpE* gene are shown in SEQ ID NO:18 and 19. The sequence of the *clpB* gene is shown in SEQ ID NO:20, the corresponding primers for the isolation of the *clpB* gene are shown in SEQ ID NO:21 and 22. In order to construct pCHAP-1 the PCR fragment containing the *dnaKJ* operon was recleaved with *SmaI* and *BamHI* and, as described above, ligated into a vector fragment of pKKT5 which had also been linearized with *SmaI* and *BamHI*. pCHAP-2 was constructed by means of a three-fold ligation with the

*EcoRI-PstI* fragment of the *grpE* gene, the *PstI-HindIII* fragment of the *clpB*-gene and a vector fragment of pKKT5 linearized with *EcoRI* and *HindIII*. p-CHAP-3 in which the *clpB* gene is present alone as a transcription unit, is derived from pCHAP-2 by ligating the *PstI-HindIII* fragment from pCHAP-2 into the vector fragment of pKKT5 linearized with *EcoRI* and *HindIII* as described above. Before the ligation reaction the non-compatible ends of the two fragments were prepared with Klenow polymerase (Roche Diagnostics) according to the manufacturer's instructions. pCHAP-4 is a pCHAP-1 derivative whose insert was extended by the *grpE* gene from pCHAP-2 and thus the artificial transcription unit comprises the genes for DnaK, DnaJ and GrpE. As a result of the Shine Dalgarno sequence which is suboptimal in this case, the expression of *grpE* should be reduced compared to pCHAP-2 and thus be better adapted to the expression of *dnaKJ* (Diamant & Goloubinoff, 1998; Pierpaoli et al., 1998). In order to construct pCHAP-4 the *EcoRI-AvaI* fragment from pCHAP-2 was inserted into the *BamHI* cleavage site of pCHAP-1 after the non-compatible ends of the two fragments had been prepared with Klenow polymerase (Roche Diagnostics) according to the manufacturer's instructions. The final construct p-CHAP-5 is a pCHAP-4 derivative which contains the insert of pCHAP-3 as additional genetic information. For this the *BspLU11I-NdeI* fragment in pCHAP-4 was replaced by the *BspLU11I-SspI* fragment from pCHAP-3 by restriction and ligation as already described several times. In order to ensure the compatibility of the ends, the overhanging ends generated by *NdeI* were previously filled in with Klenow polymerase (Roche Diagnostics) according to the manufacturer's instructions.

The effect of combining the expression plasmid pAMV $\alpha\beta$ -4

with the various helper plasmids pCHAP-1 to -5 on the overproduction of active AMV-RT was examined. At least under the modified standard expression conditions all helper plasmids considerably increased the previous yields of active AMV-RT and as expected the helper plasmid pCHAP-5 gave the best result. This was confirmed by SDS-PAGE as well as by subsequent purification and activity determination.

## **5. Example:**

### **Analytical methods**

#### **5.1. Test for reverse transcriptase activity (test A)**

During the purification, the reverse transcriptase activity in the fractions was detected by means of a non-radioactive test system. The "reverse transcriptase assay non-radioactive" (Roche Molecular Biochemicals, cat. No. 1468120) was used for this. The incubation period was shortened to 30 minutes.

#### **5.2. Test for reverse transcriptase activity (test B)**

The specific reverse transcriptase activity of the pools was determined by a radioactive test system. Reverse transcriptase activity was determined in a test volume of 100  $\mu$ l (50 mM Tris/HCl, pH 8.3 (37°C), 40 mM KCl, 6 mM MgCl<sub>2</sub>, 0.5 mM dTTP, 0.04 OD<sub>260</sub> nm poly (A) x dT<sub>15</sub>, 0.1  $\mu$ M [3H]-dTTP). AMV-RT (5  $\mu$ l) was added in suitable dilutions. After incubating for 10 min at 37°C, the reaction was stopped with 10 % TCA solution (500  $\mu$ l). The radioactively-labelled product that formed was washed on a nitrocellulose filter after precipitation. The incorporation rate of radioactivity was measured in

a scintillation counter and the RT activity of the sample was calculated. One enzyme unit was defined as the amount of AMV-RT which incorporated 1.0 nMol TMP into acid insoluble product in 10 min at 37°C.

### 5.3. Test for DNA polymerase

The activity of DNA polymerase from *E. coli* was determined by measuring the nick translation. The DNA polymerase was detected by means of a non-radioactive nick translation test. The nick translation was carried out in a test volume of 50  $\mu$ l (50 mM Tris/HCl, pH 7.5, 10 mM MgCl<sub>2</sub>, 0.1 mM DTE, 28.875  $\mu$ M DIG-dUTP, 1.444  $\mu$ M bio-16-dUTP, 95.865  $\mu$ M dTTP, 20  $\mu$ M dATP, 20  $\mu$ M dCTP, 20  $\mu$ M dGTP, 1  $\mu$ g pBR322, 1 pg DNaseI). After adding the samples (1  $\mu$ l) the reaction mixture was incubated for 30 min at 37°C. Afterwards the reaction mixture was transferred to streptavidin-coated microtitre plates. Subsequent treatment and evaluation of the test was carried out analogously to the "reverse transcriptase assay, non-radioactive" (Roche Molecular Biochemicals, Cat. No. 1468120).

### 5.4 Test for contaminating activities

The test for the presence of contaminating foreign activities was carried out in a solution composed of 10 mM Tris/HCl, pH 7.5, 10 mM MgCl<sub>2</sub>, 1 mM DTE.

Suitable samples of the individual enzyme fractions were incubated with the corresponding nucleic acids. So-called nicking activity was detected by incubation with the plasmid pBR322 (1  $\mu$ g) for 2-16 hours at 37°C. Unspecific nucleases were detected by incubation with



lambda-DNA/*Eco*RI, *Hind*III (1 µg) for 2-16 hours at 37°C. Unspecific RNases were detected by incubation for 2-4 hours at 37°C with MSII-RNA (5 µg).

For the test for contamination by exonucleases, the samples were incubated for 4 hours at 37°C with 4 µg [3H]-labelled DNA and afterwards the released [3H]-labelled nucleotides were determined.

## 6. Examples:

### Purification and functional test

#### 6.1. AMV-RT from *E. coli* LE392 pAMV- $\alpha_{lacIq-Arg}$ /pAMV- $\beta_{dnaY-His}$ construct

##### 6.1.1. Purification

*E. coli* cells which overexpress both chains of the AMV-RT (see above) were used as the starting material to purify the recombinant AMV-RT.

The AMV-RT was purified at 4°C. The purification was carried out by chromatographic methods after cell lysis and separation of the nucleic acids. The purification process yields a recombinant AMV-RT which is free of contaminating enzyme activities and in RT-PCR has the same functionality as an AMV-RT purified from native material.

#### Buffer:

buffer A: 50 mM Tris/HCl, pH 7.9, 0.5 M KCl, 0.02 % Triton X-100, 20 % glycerol,

buffer B: 20 mM Tris/HCl, pH 7.9, 0.25 M KCl, 0.02 %

Triton X-100, 10 % glycerol,  
buffer C: 20 mM Tris/HCl, pH 7.9, 0.25 M KCl, 0.02 %  
Triton X-100, 10 % glycerol, 1 M imidazole,  
buffer D: 50 mM Tris/HCl, pH 8.2, 0.1 mM EDTA, 1 mM DTT,  
0.02 % Triton X-100, 10 % glycerol,  
buffer E: 20 mM potassium phosphate, pH 7.1, 0.1 mM  
EDTA, 1 mM DTT, 0.02 % Triton X-100, 10 % glycerol,  
storage buffer: 200 mM potassium phosphate, pH 7.2, 2 mM  
DTT, 0.2 % Triton X-100, 50 % glycerol.

*Cell lysis:*

200 ml buffer A was added to ca. 50 g *E. coli* LE392  
cells (pAMV- $\alpha_{lacIq}$ -Arg/pAMV- $\beta_{dnaX}$ -His) which were thawed  
and suspended. Two tablets of Complete (Roche Molecular  
Biochemicals, cat. No. 1697498) were added to the  
suspension. Subsequently the cells were lysed by means  
of ultrasound (Branson sonicator) while cooling  
(temperature: < 10°C). The degree of lysis of the cell  
suspension that was achieved was typically 40-50 %.

*Precipitation of nucleic acids:*

Afterwards the nucleic acids were removed by means of  
polymin precipitation. 5 ml of a 10 % polymin P solution  
was added dropwise. If the precipitation was incomplete,  
further dropwise addition was carried out. After  
incubation for 30 min at 4°C, centrifugation was carried  
out (30 min, 13000 rpm, 4°C).

*Chromatographic purifications:*

*Affinity chromatography on a Ni-chelate column:*

The clear centrifugation supernatant was diluted with  
buffer B (1+1) and absorbed to a nickel-loaded chelating  
sepharose ff column (2.6 cm x 10 cm, Pharmacia) which

had been equilibrated with buffer B, it was then washed with ca. 500 ml buffer B, afterwards with 200 ml buffer B + 10 mM imidazole and 200 ml buffer B + 20 mM imidazole. The enzyme was eluted with a linear gradient of buffer B + 20 mM imidazole and buffer C in a total volume of 500 ml. The flow rate was 5 ml per minute, the fraction size was 20 ml per fraction. The enzyme eluted between 50 mM and 200 mM imidazole. The pool of active fractions was dialysed against buffer D.

*Chromatography on heparin-sepharose:*

The dialysed pool was subsequently absorbed to a heparin-sepharose ff column equilibrated with buffer D (1.6 cm x 10 cm, Pharmacia) and washed with ca. 200 ml buffer D, then with ca. 200 ml buffer D + 300 mM KCl. The enzyme was eluted with a linear gradient of buffer D + 300 mM KCl and buffer D + 1 M KCl in a total volume of 200 ml. The flow rate was 2.5 ml per min, the fraction size was 10 ml. The AMV-RT eluted at a KCl concentration of 500 mM to 700 mM.

*Chromatography on S-sepharose ff:*

The RT-active fractions were pooled and dialysed against buffer E. The dialysate was loaded onto a S-sepharose ff column equilibrated with buffer E (1.6 cm x 10 cm, Pharmacia). After washing with ca. 200 ml buffer E, the enzyme was eluted with a linear gradient of buffer E and buffer E + 1 M KCl in a total volume of 400 ml. The flow rate was 2.5 ml per minute, the fraction size was 10 ml.

*Chromatography on hydroxylapatite:*

The RT-active fractions were pooled and dialysed against buffer E. The dialysate was loaded onto a HA-ultrogel column equilibrated with buffer E (1.6 cm x 10 cm,

Biosepra). After washing with ca. 200 ml buffer E, the enzyme was eluted with a linear gradient of buffer E and buffer E + 0.5 M potassium phosphate in a total volume of 400 ml. The flow rate was 2.5 ml per minute, the fraction size was 10 ml.

The RT-active fractions were pooled and dialysed against storage buffer. Application buffer containing SDS and  $\beta$ -mercaptoethanol was added to the purified protein and the sample was denatured by boiling (5 min, 100°C). Subsequently 20  $\mu$ l aliquots were analysed by an analytical SDS gel (4-20 %) (Laemmli UK., 1970, Nature **227**: 555-557). The  $\alpha$ - and  $\beta$ -subunits of AMV-RT were found in equimolar ratios.

The described method yields a stable AMV-RT with an equimolar distribution of the  $\alpha$ - and  $\beta$ -subunits. The enzyme obtained is functional in RT-PCR.

#### 6.1.2. Functional test in RT-PCR

The recombinant AMV reverse transcriptase that was obtained was examined in a functional test. The functional test consisted of a reverse transcription (RT) coupled with a polymerase chain reaction (PCR). For this 5 units of the recombinant AMV reverse transcriptase was used like the enzyme mixture of the Titan TM One Tube PCR System (cat. No. 1888382, Roche Molecular Biochemicals). A 1.8 kb fragment of the human dystrophin gene was amplified. 10 ng human muscle RNA was used as a template. The primers (400 nM) were the Dys primer 2rev (5'-GAG TGA ATA CAG TTT GCC CAT GGA TTG-3') and the Dys primer 8for (5'-AAG AAG TAG AGG ACT GTT ATG AAA GAG AAG-3'). The target was amplified in a RT-

PCR using the following program: 50°C for 30 min, 94°C for 2 min followed by 10 cycles (94°C for 10 sec, 58°C for 30 sec, 68°C for 1 min 10 sec) and 20 cycles (94°C for 10 sec, 58°C for 30 sec, 68°C for 1 min 10 sec; + 10 sec/cycle). Subsequently it was incubated for 7 min at 68°C. The reaction products of the RT-PCR were separated after stopping the reaction on a 1 % agarose gel (fig. 1).

Figure 1 shows the amplification product of the RT-PCR having a size of 1.8 kb which was obtained using native purified AMV-RT (lane 2) and AMV-RT that was obtained by recombinant means (lane 3). Lanes 1 and 4 show a DNA molecular weight marker VI (cat. No. 1062590, Roche Molecular Biochemicals).

## **6.2 AMV-RT from *E. coli* LE392 pAMV $\alpha\beta$ -4 + pCHAP-5 construct**

### **6.2.1 Purification**

*E. coli* LE392 pAMV $\alpha\beta$ -4 + pCHAP-5 cells which overexpress both chains of the AMV-RT (see above) were used as the starting material to purify the recombinant AMV-RT.

The AMV-RT was purified at 4°C. The purification was carried out by chromatographic methods after cell lysis and separation of the nucleic acids. The purification yields a recombinant AMV-RT which is free of contaminating enzyme activities and in RT-PCR has the same functionality as an AMV-RT purified from native material.

*Buffer:*

buffer A: 50 mM NaPO<sub>4</sub>, pH 7.2, 1 M NaCl, 3 mM 2-mercapto-ethanol, 10 % glycerol,

buffer B: 50 mM NaPO<sub>4</sub>, pH 5.0, 1 M NaCl, 3 mM 2-mercapto-ethanol, 10 % glycerol,

buffer C: 50 mM NaPO<sub>4</sub>, pH 6.0, 1 M NaCl, 3 mM 2-mercapto-ethanol, 10 % glycerol, 0.2 M imidazole,

buffer D: 50 mM NaPO<sub>4</sub>, pH 7.7, 1 M NaCl, 3 mM 2-mercapto-ethanol, 10 % glycerol, 0.5 M imidazole

buffer E: 50 mM NaPO<sub>4</sub>, pH 6.0, 3 mM 2-mercaptoethanol, 10 % glycerol,

storage buffer: 200 mM potassium phosphate, pH 7.2, 2 mM DTT, 0.2 % Triton X-100, 50 % glycerol.

*Cell lysis:*

Ca. 50 g *E coli* LE392 pAMV- $\alpha\beta$ -4+pCHAP-5 cells were mixed with 400 ml buffer A, thawed and suspended. Two tablets of Complete (Roche Molecular Biochemicals, cat. No. 1697498) were added to the suspension. Subsequently the cells were lysed by means of ultrasound (Branson sonicator) while cooling (temperature: < 10°C). The degree of lysis of the cell suspension that was achieved was typically 40-50 %.

*Precipitation of nucleic acids:*

Afterwards the nucleic acids were removed by means of polymin precipitation. 5 ml of a 10 % polymin P solution was added dropwise. If the precipitation was incomplete, further dropwise addition was carried out. After incubation for 30 min at 4°C, centrifugation was carried out (30 min, 13000 rpm, 4°C).

*Chromatographic purifications:*

*Affinity chromatography on a Ni-chelate column:*

The clear centrifugation supernatant was absorbed to a nickel-loaded chelating sepharose ff column (2.6 cm x 10 cm, Pharmacia) which had been equilibrated with buffer A, it was then washed with ca. 500 ml buffer A, afterwards with 500 ml buffer B and 500 ml buffer C. The enzyme was eluted with buffer D in a total volume of 500 ml. The flow rate was 5 ml per minute, the fraction size was 20 ml per fraction. The pool of active fractions was dialysed against buffer E.

*Chromatography on heparin-sepharose:*

The dialysed pool was subsequently absorbed to a heparin-sepharose ff column (1.6 cm x 10 cm, Pharmacia) equilibrated with buffer E + 250 mM NaCl and washed with ca. 200 ml buffer E + 250 mM NaCl. The enzyme was eluted with a linear gradient of buffer E + 250 mM NaCl and buffer E + 1 M NaCl in a total volume of 200 ml. The flow rate was 2.5 ml per min, the fraction size was 10 ml. The AMV-RT eluted at an NaCl concentration of 500 mM to 700 mM.

The RT-active fractions were pooled and dialysed against storage buffer. Application buffer containing SDS and  $\beta$ -mercaptoethanol was added to the purified protein and the sample was denatured by boiling (5 min, 100°C). Subsequently 20  $\mu$ l aliquots were analysed by an analytical SDS gel (4-20 %) (Laemmli UK., 1970, Nature **227**: 555-557). The  $\alpha$ - and  $\beta$ -subunits of AMV-RT were found in equimolar ratios (fig. 2, lane 6).

Figure 2 shows an SDS gel with samples from the AMV-RT purification

lane 1: molecular weight marker

lane 2: native AMV

lane 3: cell lysis

lane 4: Ni-chelate sepharose, wash with buffer C

lane 5: Ni-chelate pool

lane 6: rec. AMV-RT final preparation

The described method yields a stable AMV-RT with an equimolar distribution of the  $\alpha$ - and  $\beta$ -subunits. The enzyme obtained is functional in RT-PCR.

#### 6.2.2. Functional test in RT-PCR

The recombinant AMV reverse transcriptase that was obtained was examined in a functional test. The functional test consists of a reverse transcription (RT), followed by a polymerase chain reaction (PCR). 10 units of the recombinant AMV reverse transcriptase was used for this. A 8 kb, 10 kb, 12 kb and a 13.5 kb fragment of the human dystrophin gene was amplified.

1  $\mu$ g human muscle RNA was used as a template. The primers (400 nM) were the Dys primer 2 for (5'-CAA TCC ATG GGC AAA CTG TAT TCA CTC-3') and Dys primer 5 rev (5'-CGT CCC GTA TCA TAA ACA TTC AGC AGC-3') for 8 kb, Dys primer 8 for (5'-AAG AAG TAG AGG ACT GTT ATG AAA GAG AA-3') and 5 rev for 10 kb, Dys primer 8 for and Dys primer 9 rev (5'-AGC AGG TAA GCC TGG ATG ACT GAC TAG AAG-3') for 12 kb and Dys primer 8 for and 10 rev (5'-AAT CAA TCA ACC AAC CGA AAT CTC ACT CTG-3') for 13.5 kb. The cDNA synthesis was carried out for 60 min at 42°C.



The cDNA synthesis was carried out according to the instructions in the product information for the AMV reverse transcriptase (cat. No. 1495062, Roche Molecular Biochemicals).

The Expand Long Template PCR System (cat. No. 1681834, Roche Molecular Biochemicals) was used for the PCR. The target was amplified using the following PCR program: 94°C for 2 min, followed by 10 cycles (94°C for 10 sec, 60°C for 30 sec, 68°C for 10 min) and 20 cycles (94°C for 10 sec, 60°C for 30 sec, 68°C for 10 min 10 + 10 sec/cycle). Subsequently it was incubated for 5 min at 68°C. After stopping the reaction, the reaction products of the RT-PCR were separated on 1 % agarose gel (fig. 3). Lanes 3 and 6 show a DNA molecular weight marker X (cat. no. 1498037, Roche Molecular Biochemicals).

Figure 3 shows an agarose gel on which the reaction products of the RT-PCR using recombinant AMV-RT were separated; lane 1: 8 kb amplification product, lane 2: 10 kb amplification product, lane 3: DNA length standard X, lane 4: 12 kb amplification product, lane 5: 13.5 kb amplification product, lane 6: DNA length standard X.

SEQUENCE LISTING

<110> Roche Diagnostics GmbH

<120> Method for producing an active heterodimeric AMV-RT in prokaryotic cells

<130> 5272/00/DE

<140>

<141>

<160> 22

<170> PatentIn Ver. 2.1

<210> 1

<211> 38

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 1

gatgactgga attcatgact gttgcgctac atctggct

38

<210> 2

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 2

gatgactgct gcagttatta tgcaaaaaga gggctcgcct

40

<210> 3

<211> 41

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 3

gatgactgct gcagttatta atacgcttga aaggtggctt g

41

<210> 4

<211> 1716

<212> DNA

<213> Avian Myeloblastosis Virus

<400> 4

actgtttgctgac tacatctggc tattccgctc aaatggaagc caaaccacac gcctgtgtgg 60  
attgaccagt ggcccttcc tgaaggtaaa cttgtagcgc taacgcaatt agtggaaaaa 120  
gaattacagt taggacatat agaaccttca cttagtgtgt ggaacacacc tgtctttgtg 180  
atccggaagg cttccgggtc ttatcgctta ttgcatgact tgcgcgctgt taacgctaag 240  
cttgttcctt ttggggccgt ccaacagggg gcgcgggttc tctccgctgt cccgcgtggc 300  
tggccctga tggctctaga cctcaaggat tgcttctttt ctattcctct tgcggaacaa 360  
gatcgcaag cttttgcatt tacgctcccc tctgtgaata accaggcccc cgctcgaaga 420  
ttccaatgga aggtcttgcc ccaagggatg acctgttctc ccactatctg tcagttgata 480

gtgggtcaaa	tacttgagcc	cttgcgactc	aagcacccat	ctctgcgcac	gttgccattat	540
atggatgac	ttttgctagc	cgcctcaagt	catgatgggt	tggaaagcggc	aggggaggag	600
gttatcagta	catttgaaag	agccgggttc	accatttcgc	ctgataaggt	ccagaggag	660
cccgagtagc	aatatcttgg	gtacaagtta	ggcagtagct	atgtagcacc	cgtaggcctg	720
gtagcagaac	ccaggatagc	caccttggtg	gatgttcaga	agctggtggg	gtcacttcag	780
tggcttcgcc	cagcgctagg	aatcccgcc	cgactgatgg	gcccccttta	tgagcagtta	840
cgagggtcag	atcctaacga	ggcgagggaa	tggaaatctag	acatgaaaat	ggcctggaga	900
gagatcgtag	agctcagcac	cactgctgcc	ttggaacgat	gggaccctgc	cctgcctctg	960
gaaggagcgg	tcgctagatg	tgaacagggg	gcaatagggg	tcttgggaca	gggactgtcc	1020
acacacccaa	ggccatgttt	gtgggttattc	tccacccaac	ccaccaaggc	gtttactgct	1080
tgggttagaag	tgctcaacct	tttgattact	aagctacgtg	cttcggcagt	gcgaaccttt	1140
ggcaaggagg	ttgatatact	cctgttgcc	gcatgctttc	gggaggacct	tccgctcccc	1200
gaggggatcc	tgtagccct	taggggggtt	gcaggaaaaa	tcaggagtag	tgacacgcca	1260
tctatttttg	acattgcgcg	tccactgcac	gtttctctga	aagtgagggt	taccgaccac	1320
cctgtaccgg	gacccactgt	ctttaccgac	gcctcctcaa	gcaccataa	gggggtggta	1380
gtctggaggg	agggcccaag	gtgggagata	aaagaaatag	ctgatttggtg	ggcaagtgtg	1440
caacaactgg	aagcaogcgc	tgtggccatg	gcacttctgc	tgtggccgac	aacgcccact	1500
aatgtagtga	ctgactctgc	gtttgttgcg	aaaatgttac	tcaagatggg	gcaggaggga	1560
gtcccgctca	cagcggcgcg	ttttatttta	gaggatgcgt	taagccaaag	gtcagccatg	1620
gccgcggttc	tccacgtgcg	gagtcattct	gaggtgccag	ggtttttcac	agaaggaaat	1680
gacgtggcag	atagccaagc	cacctttcaa	gcgtat			1716

<210> 5

<211> 2574

<212> DNA

<213> Avian Myeloblastosis Virus

<400> 5

actgttgccg	tacatctggc	tattccgctc	aaatggaagc	caaaccacac	gcctgtgtgg	60
attgaccagt	ggcccccttc	tgaaggtaaa	cttgtagcgc	taacgcaatt	agtggaaaaa	120
gaattacagt	taggacatat	agaacottca	cttagttgct	ggaacacacc	tgtctttgtg	180
atccggaagg	cttcggggtc	ttatcgctta	ttgcatgact	tgcgcgctgt	taacgctaag	240
cttggttcct	ttggggcggt	ccaacagggg	gcgcgggttc	tctccgcgct	cccgcgtggc	300
tggcccccta	tggtcctaga	cctcaaggat	tgcttctttt	ctattcctct	tgcggaacaa	360
gatcgcgagg	cttttgcaat	tacgtctccc	tctgtgaata	accaggcccc	cgctcgaaga	420
ttccaatgga	aggtcttgcc	ccaagggatg	acctgttctc	ccactatctg	tcagttgata	480
gtgggtcaaa	tacttgagcc	cttgcgactc	aagcacccat	ctctgcgcac	gttgccattat	540
atggatgac	ttttgctagc	cgcctcaagt	catgatgggt	tggaaagcggc	aggggaggag	600
gttatcagta	catttgaaag	agccgggttc	accatttcgc	ctgataaggt	ccagaggag	660
cccgagtagc	aatatcttgg	gtacaagtta	ggcagtagct	atgtagcacc	cgtaggcctg	720
gtagcagaac	ccaggatagc	caccttggtg	gatgttcaga	agctggtggg	gtcacttcag	780
tggcttcgcc	cagcgctagg	aatcccgcc	cgactgatgg	gcccccttta	tgagcagtta	840
cgagggtcag	atcctaacga	ggcgagggaa	tggaaatctag	acatgaaaat	ggcctggaga	900
gagatcgtag	agctcagcac	cactgctgcc	ttggaacgat	gggaccctgc	cctgcctctg	960
gaaggagcgg	tcgctagatg	tgaacagggg	gcaatagggg	tcttgggaca	gggactgtcc	1020
acacacccaa	ggccatgttt	gtgggttattc	tccacccaac	ccaccaaggc	gtttactgct	1080
tgggttagaag	tgctcaacct	tttgattact	aagctacgtg	cttcggcagt	gcgaaccttt	1140
ggcaaggagg	ttgatatact	cctgttgcc	gcatgctttc	gggaggacct	tccgctcccc	1200
gaggggatcc	tgtagccct	taggggggtt	gcaggaaaaa	tcaggagtag	tgacacgcca	1260
tctatttttg	acattgcgcg	tccactgcac	gtttctctga	aagtgagggt	taccgaccac	1320
cctgtaccgg	gacccactgt	ctttaccgac	gcctcctcaa	gcaccataa	gggggtggta	1380
gtctggaggg	agggcccaag	gtgggagata	aaagaaatag	ctgatttggtg	ggcaagtgtg	1440
caacaactgg	aagcacgcgc	tgtggccatg	gcacttctgc	tgtggccgac	aacgcccact	1500
aatgtagtga	ctgactctgc	gtttgttgcg	aaaatgttac	tcaagatggg	gcaggaggga	1560
gtcccgctca	cagcggcgcg	ttttatttta	gaggatgcgt	taagccaaag	gtcagccatg	1620
gccgcggttc	tccacgtgcg	gagtcattct	gaagtgcag	ggtttttcac	agaaggaaat	1680
gacgtggcag	atagccaagc	cacctttcaa	gcgtatccct	tgagagaggc	taaagatctc	1740
cataccgctc	tccatatcgg	accccgcgcg	ctatccaaag	cgtgtaatat	atctatgcag	1800
caggctaggg	aggttggtca	gacctgcccc	cattgttaatt	cagccccctgc	gttgaggaggc	1860
ggggtaaaac	ctagggggtt	gggaccctca	cagatatggc	agacagactt	tacactagag	1920
cctagaaatg	ctccccgttc	ctggctcgct	gttactgtgg	ataccgcctc	atctgcgata	1980
gtcgtaaactc	agcatggccg	tgtcacatcg	gttgctgcac	aacatcattg	ggccacggct	2040
atcgccgttt	tgggaagacc	aaaggccata	aaaacagata	atgggtcctg	cttcacgtct	2100

```

aatccacgc gagagtggct cgcgagatgg gggatagcac acaccaccgg gattccgggt 2160
aattcccagg gtcaagctat ggtagagcgg gccaacccgg tcctgaaaga taagatccgt 2220
gtgcttgccg agggggatgg ctttatgaaa agaatcccca ccagcaaaca gggggaacta 2280
ttagccaagg caatgtatgc ccttaatcac tttagcgtg gtgaaaacac aaaaacaccg 2340
atacaaaaac actggagacc taccgttctt acagaaggac ccccggttaa aatacgaata 2400
gagacagggg agtgggaaaa aggatggaac gtgctggtct ggggacgagg ttatgcagct 2460
gtgaaaaaca gggacactga taagggtatt tgggtaccct ctcgaaaagt taaaccggac 2520
atcgcccaaa aggatgaggt gactaagaaa gatgaggcga gccctctttt tgca 2574

```

<210> 6

<211> 572

<212> PRT

<213> Avian Myeloblastosis Virus

<400> 6

Thr Val Ala Leu His Leu Ala Ile Pro Leu Lys Trp Lys Pro Asn His  
1 5 10 15

Thr Pro Val Trp Ile Asp Gln Trp Pro Leu Pro Glu Gly Lys Leu Val  
20 25 30

Ala Leu Thr Gln Leu Val Glu Lys Glu Leu Gln Leu Gly His Ile Glu  
35 40 45

Pro Ser Leu Ser Cys Trp Asn Thr Pro Val Phe Val Ile Arg Lys Ala  
50 55 60

Ser Gly Ser Tyr Arg Leu Leu His Asp Leu Arg Ala Val Asn Ala Lys  
65 70 75 80

Leu Val Pro Phe Gly Ala Val Gln Gln Gly Ala Pro Val Leu Ser Ala  
85 90 95

Leu Pro Arg Gly Trp Pro Leu Met Val Leu Asp Leu Lys Asp Cys Phe  
100 105 110

Phe Ser Ile Pro Leu Ala Glu Gln Asp Arg Glu Ala Phe Ala Phe Thr  
115 120 125

Leu Pro Ser Val Asn Asn Gln Ala Pro Ala Arg Arg Phe Gln Trp Lys  
130 135 140

Val Leu Pro Gln Gly Met Thr Cys Ser Pro Thr Ile Cys Gln Leu Ile  
145 150 155 160

Val Gly Gln Ile Leu Glu Pro Leu Arg Leu Lys His Pro Ser Leu Arg  
165 170 175

Met Leu His Tyr Met Asp Asp Leu Leu Leu Ala Ala Ser Ser His Asp  
180 185 190

Gly Leu Glu Ala Ala Gly Glu Glu Val Ile Ser Thr Leu Glu Arg Ala  
195 200 205

Gly Phe Thr Ile Ser Pro Asp Lys Val Gln Arg Glu Pro Gly Val Gln  
210 215 220

Tyr Leu Gly Tyr Lys Leu Gly Ser Thr Tyr Val Ala Pro Val Gly Leu  
225 230 235 240

Val Ala Glu Pro Arg Ile Ala Thr Leu Trp Asp Val Gln Lys Leu Val  
245 250 255

Gly	Ser	Leu	Gln	Trp	Leu	Arg	Pro	Ala	Leu	Gly	Ile	Pro	Pro	Arg	Leu
			260					265					270		
Met	Gly	Pro	Phe	Tyr	Glu	Gln	Leu	Arg	Gly	Ser	Asp	Pro	Asn	Glu	Ala
		275					280					285			
Arg	Glu	Trp	Asn	Leu	Asp	Met	Lys	Met	Ala	Trp	Arg	Glu	Ile	Val	Gln
	290					295					300				
Leu	Ser	Thr	Thr	Ala	Ala	Leu	Glu	Arg	Trp	Asp	Pro	Ala	Leu	Pro	Leu
305					310					315					320
Glu	Gly	Ala	Val	Ala	Arg	Cys	Glu	Gln	Gly	Ala	Ile	Gly	Val	Leu	Gly
				325					330					335	
Gln	Gly	Leu	Ser	Thr	His	Pro	Arg	Pro	Cys	Leu	Trp	Leu	Phe	Ser	Thr
			340					345					350		
Gln	Pro	Thr	Lys	Ala	Phe	Thr	Ala	Trp	Leu	Glu	Val	Leu	Thr	Leu	Leu
		355					360					365			
Ile	Thr	Lys	Leu	Arg	Ala	Ser	Ala	Val	Arg	Thr	Phe	Gly	Lys	Glu	Val
	370					375					380				
Asp	Ile	Leu	Leu	Leu	Pro	Ala	Cys	Phe	Arg	Glu	Asp	Leu	Pro	Leu	Pro
385					390					395					400
Glu	Gly	Ile	Leu	Leu	Ala	Leu	Arg	Gly	Phe	Ala	Gly	Lys	Ile	Arg	Ser
				405					410					415	
Ser	Asp	Thr	Pro	Ser	Ile	Phe	Asp	Ile	Ala	Arg	Pro	Leu	His	Val	Ser
			420					425					430		
Leu	Lys	Val	Arg	Val	Thr	Asp	His	Pro	Val	Pro	Gly	Pro	Thr	Val	Phe
		435				440					445				
Thr	Asp	Ala	Ser	Ser	Ser	Thr	His	Lys	Gly	Val	Val	Val	Trp	Arg	Glu
	450					455					460				
Gly	Pro	Arg	Trp	Glu	Ile	Lys	Glu	Ile	Ala	Asp	Leu	Gly	Ala	Ser	Val
465					470					475					480
Gln	Gln	Leu	Glu	Ala	Arg	Ala	Val	Ala	Met	Ala	Leu	Leu	Leu	Trp	Pro
				485					490					495	
Thr	Thr	Pro	Thr	Asn	Val	Val	Thr	Asp	Ser	Ala	Phe	Val	Ala	Lys	Met
			500					505					510		
Leu	Leu	Lys	Met	Gly	Gln	Glu	Gly	Val	Pro	Ser	Thr	Ala	Ala	Ala	Phe
		515					520					525			
Ile	Leu	Glu	Asp	Ala	Leu	Ser	Gln	Arg	Ser	Ala	Met	Ala	Ala	Val	Leu
	530					535					540				
His	Val	Arg	Ser	His	Ser	Glu	Val	Pro	Gly	Phe	Phe	Thr	Glu	Gly	Asn
545					550					555					560
Asp	Val	Ala	Asp	Ser	Gln	Ala	Thr	Phe	Gln	Ala	Tyr				
				565					570						

<210> 7  
 <211> 858  
 <212> PRT  
 <213> Avian Myeloblastosis Virus

<400> 7  
 Thr Val Ala Leu His Leu Ala Ile Pro Leu Lys Trp Lys Pro Asn His  
 1 5 10 15  
 Thr Pro Val Trp Ile Asp Gln Trp Pro Leu Pro Glu Gly Lys Leu Val  
 20 25 30  
 Ala Leu Thr Gln Leu Val Glu Lys Glu Leu Gln Leu Gly His Ile Glu  
 35 40 45  
 Pro Ser Leu Ser Cys Trp Asn Thr Pro Val Phe Val Ile Arg Lys Ala  
 50 55 60  
 Ser Gly Ser Tyr Arg Leu Leu His Asp Leu Arg Ala Val Asn Ala Lys  
 65 70 75 80  
 Leu Val Pro Phe Gly Ala Val Gln Gln Gly Ala Pro Val Leu Ser Ala  
 85 90 95  
 Leu Pro Arg Gly Trp Pro Leu Met Val Leu Asp Leu Lys Asp Cys Phe  
 100 105 110  
 Phe Ser Ile Pro Leu Ala Glu Gln Asp Arg Glu Ala Phe Ala Phe Thr  
 115 120 125  
 Leu Pro Ser Val Asn Asn Gln Ala Pro Ala Arg Arg Phe Gln Trp Lys  
 130 135 140  
 Val Leu Pro Gln Gly Met Thr Cys Ser Pro Thr Ile Cys Gln Leu Ile  
 145 150 155 160  
 Val Gly Gln Ile Leu Glu Pro Leu Arg Leu Lys His Pro Ser Leu Arg  
 165 170 175  
 Met Leu His Tyr Met Asp Asp Leu Leu Leu Ala Ala Ser Ser His Asp  
 180 185 190  
 Gly Leu Glu Ala Ala Gly Glu Glu Val Ile Ser Thr Leu Glu Arg Ala  
 195 200 205  
 Gly Phe Thr Ile Ser Pro Asp Lys Val Gln Arg Glu Pro Gly Val Gln  
 210 215 220  
 Tyr Leu Gly Tyr Lys Leu Gly Ser Thr Tyr Val Ala Pro Val Gly Leu  
 225 230 235 240  
 Val Ala Glu Pro Arg Ile Ala Thr Leu Trp Asp Val Gln Lys Leu Val  
 245 250 255  
 Gly Ser Leu Gln Trp Leu Arg Pro Ala Leu Gly Ile Pro Pro Arg Leu  
 260 265 270  
 Met Gly Pro Phe Tyr Glu Gln Leu Arg Gly Ser Asp Pro Asn Glu Ala  
 275 280 285  
 Arg Glu Trp Asn Leu Asp Met Lys Met Ala Trp Arg Glu Ile Val Gln  
 290 295 300

Leu Ser Thr Thr Ala Ala Leu Glu Arg Trp Asp Pro Ala Leu Pro Leu  
 305 310 315 320  
 Glu Gly Ala Val Ala Arg Cys Glu Gln Gly Ala Ile Gly Val Leu Gly  
 325 330 335  
 Gln Gly Leu Ser Thr His Pro Arg Pro Cys Leu Trp Leu Phe Ser Thr  
 340 345 350  
 Gln Pro Thr Lys Ala Phe Thr Ala Trp Leu Glu Val Leu Thr Leu Leu  
 355 360 365  
 Ile Thr Lys Leu Arg Ala Ser Ala Val Arg Thr Phe Gly Lys Glu Val  
 370 375 380  
 Asp Ile Leu Leu Leu Pro Ala Cys Phe Arg Glu Asp Leu Pro Leu Pro  
 385 390 395 400  
 Glu Gly Ile Leu Leu Ala Leu Arg Gly Phe Ala Gly Lys Ile Arg Ser  
 405 410 415  
 Ser Asp Thr Pro Ser Ile Phe Asp Ile Ala Arg Pro Leu His Val Ser  
 420 425 430  
 Leu Lys Val Arg Val Thr Asp His Pro Val Pro Gly Pro Thr Val Phe  
 435 440 445  
 Thr Asp Ala Ser Ser Ser Thr His Lys Gly Val Val Val Trp Arg Glu  
 450 455 460  
 Gly Pro Arg Trp Glu Ile Lys Glu Ile Ala Asp Leu Gly Ala Ser Val  
 465 470 475 480  
 Gln Gln Leu Glu Ala Arg Ala Val Ala Met Ala Leu Leu Leu Trp Pro  
 485 490 495  
 Thr Thr Pro Thr Asn Val Val Thr Asp Ser Ala Phe Val Ala Lys Met  
 500 505 510  
 Leu Leu Lys Met Gly Gln Glu Gly Val Pro Ser Thr Ala Ala Ala Phe  
 515 520 525  
 Ile Leu Glu Asp Ala Leu Ser Gln Arg Ser Ala Met Ala Ala Val Leu  
 530 535 540  
 His Val Arg Ser His Ser Glu Val Pro Gly Phe Phe Thr Glu Gly Asn  
 545 550 555 560  
 Asp Val Ala Asp Ser Gln Ala Thr Phe Gln Ala Tyr Pro Leu Arg Glu  
 565 570 575  
 Ala Lys Asp Leu His Thr Ala Leu His Ile Gly Pro Arg Ala Leu Ser  
 580 585 590  
 Lys Ala Cys Asn Ile Ser Met Gln Gln Ala Arg Glu Val Val Gln Thr  
 595 600 605  
 Cys Pro His Cys Asn Ser Ala Pro Ala Leu Glu Ala Gly Val Asn Pro  
 610 615 620  
 Arg Gly Leu Gly Pro Leu Gln Ile Trp Gln Thr Asp Phe Thr Leu Glu  
 625 630 635 640

Pro Arg Met Ala Pro Arg Ser Trp Leu Ala Val Thr Val Asp Thr Ala  
645 650 655

Ser Ser Ala Ile Val Val Thr Gln His Gly Arg Val Thr Ser Val Ala  
660 665 670

Ala Gln His His Trp Ala Thr Ala Ile Ala Val Leu Gly Arg Pro Lys  
675 680 685

Ala Ile Lys Thr Asp Asn Gly Ser Cys Phe Thr Ser Lys Ser Thr Arg  
690 695 700

Glu Trp Leu Ala Arg Trp Gly Ile Ala His Thr Thr Gly Ile Pro Gly  
705 710 715 720

Asn Ser Gln Gly Gln Ala Met Val Glu Arg Ala Asn Arg Leu Leu Lys  
725 730 735

Asp Lys Ile Arg Val Leu Ala Glu Gly Asp Gly Phe Met Lys Arg Ile  
740 745 750

Pro Thr Ser Lys Gln Gly Glu Leu Leu Ala Lys Ala Met Tyr Ala Leu  
755 760 765

Asn His Phe Glu Arg Gly Glu Asn Thr Lys Thr Pro Ile Gln Lys His  
770 775 780

Trp Arg Pro Thr Val Leu Thr Glu Gly Pro Pro Val Lys Ile Arg Ile  
785 790 795 800

Glu Thr Gly Glu Trp Glu Lys Gly Trp Asn Val Leu Val Trp Gly Arg  
805 810 815

Gly Tyr Ala Ala Val Lys Asn Arg Asp Thr Asp Lys Val Ile Trp Val  
820 825 830

Pro Ser Arg Lys Val Lys Pro Asp Ile Ala Gln Lys Asp Glu Val Thr  
835 840 845

Lys Lys Asp Glu Ala Ser Pro Leu Phe Ala  
850 855

<210> 8

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 8

gatgactgga attcatgcgt cgccgctcgcc gtcgcccgtcg cactgttgcg ctacatctgg 60  
ct 62

<210> 9

<211> 65

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer



<400> 9  
gatgactgga attcatgaga ggcagccacc atcaccatca ccatactgtt gcgctacatc 60  
tggtct 65

<210> 10  
<211> 425  
<212> DNA  
<213> Escherichia coli

<400> 10  
ctgttttggc ggatgagaga agattttcag cctgatacag attaaatcag aacgcagaag 60  
cggctctgata aaacagaatt tgcctggcgg cagtagcgcg gtgggtccac ctgaccccat 120  
gccgaactca gaagtgaaac gccgtagcgc c gatggtagt gtgggggtctc cccatgcgag 180  
agtagggaac tgccaggcat caaataaaac gaaaggctca gtcgaaagac tgggcctttc 240  
gttttatctg ttgtttgtcg gtgaacgctc tcctgagtag gacaaatccg ccgggagcgg 300  
atttgaacgt tgcgaagcaa cggcccgag ggtggcgggc aggacgcccg ccataaactg 360  
ccaggcatca aattaagcag aaggccatgc tgacggatgg cttttttgcg tttctacaaa 420  
ctctt 425

<210> 11  
<211> 31  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 11  
aaaactgcag agcagtaagc cggtcataaa a 31

<210> 12  
<211> 31  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 12  
aaaactgcag cgtgctggat gaagtgtatt a 31

<210> 13  
<211> 2155  
<212> DNA  
<213> Escherichia coli

<400> 13  
atcagaatth tttttctttt tcccccttga aggggcgaag cctcatcccc atttctctgg 60  
tcaccagccg ggaaaccacg taagctccgg cgtcacccat aacagatacg gactttctca 120  
aaggagagtt atcaatgaat attcgtccat tgcatgatcg cgtgatcgtc aagcgtaaaag 180  
aagttgaaac taaatctgct ggcggcacatg ttctgacggg ctctgcagcg gctaaatcca 240  
cccgcggcga agtgctggct gtcggcaatg gccgtatcct tgaaaatggc gaagtgaagc 300  
cgctggatgt gaaagtgggc gacatcggtt ttttcaacga tggctacggg gtgaaatctg 360  
agaagatcga caatgaagaa gtgttgatca tgtccgaaag cgacattctg gcaattgttg 420  
aagcgtaatc cgcgcacgac actgaacata cgaattttaag gaataaagat aatggcagct 480  
aaagacgtaa aattcggtaa cgacgctcgt gtgaaaatgc tgcgcggcgt aaacgtactg 540  
gcagatgcag tgaaagttaac cctcgggtcca aaaggccgta acgtagttct ggataaatct 600  
ttcgggtgcac cgaccatcac caaagatggg gtttccggtt ctctgtaaat cgaactggaa 660  
gacaagttcg aaaatatggg tgcgcagatg gtgaaagaag ttgcctctaa agcaaaccgac 720  
gctgcaggcg acggtaccac cactgcaacc gtactggctc aggctatcat cactgaaggt 780  
ctgaaagctg ttgctgcggg catgaacccg atggacctga aacgtggtat cgacaaagcg 840  
gttaccgctg cagttgaaga actgaaagcg ctgtccgtac catgctctga ctctaaagcg 900  
attgctcagg ttggtaccat ctccgctaac tccgacgaaa ccgtaggtaa actgatcgct 960

gaagcgatgg	acaaagtcgg	taaagaaggc	gttatcaccg	ttgaagacgg	taccgggtctg	1020
caggacgaac	tggacgtggt	tgaaggatatg	cagttcgacc	gtggctacct	gtctccttac	1080
ttcatcaaca	agccggaaac	tggcgagta	gaactggaaa	gcccgttcat	cctgctggct	1140
gacaagaaaa	tctccaacat	ccgcgaaatg	ctgcgggttc	tggagctgt	tgccaaagca	1200
ggcaaacgcg	tgctgatcat	cgctgaagat	gtagaaggcg	aagcgtggc	aactctggtt	1260
gttaacacca	tgcgtggcat	cgtgaaagtc	gctgcggtta	aagcaccggg	cttcggcgat	1320
cgctgtaaag	ctatgctgca	ggatatcgca	accctgactg	gcggtaccgt	gatctctgaa	1380
gagatcggtg	tggagctgga	aaaagcaacc	ctggaagacc	tgggtcaggc	taaacgtggt	1440
gtgatcaaca	aagacaccac	cactatcatc	gatggcggtg	gtgaagaagc	tgcaatccag	1500
ggccgtgttg	ctcagatccg	tcagcagatt	gaagaagcaa	cttctgacta	cgaccgtgaa	1560
aaactgcagg	aacgcgtagc	gaaactggca	ggcggcggtg	cagttatcaa	agtgggtgct	1620
gctaccgaa	ttgaaatgaa	agagaaaaaa	gcacgcgttg	aagatgccct	gcacgcgacc	1680
cgtgctgcgg	tagaagaagg	cgtggttgct	ggtggtggtg	ttgcgctgat	ccgcgtagcg	1740
tctaaactgg	ctgacctgcg	tggtcagaac	gaagaccaga	acgtgggtat	caaagtgcga	1800
ctgcgtgcaa	tggagctcc	gctgcgtcag	atcgatttga	actgcggcga	agaaccgtct	1860
gttggttgcta	acaccgttaa	aggcggcgac	ggcaactacg	gttacaacgc	agcaaccgaa	1920
gaatacggca	acatgatcga	ctgggtatc	ctggatccaa	ccaaagtaac	tcgttctgct	1980
ctgcagtagc	cagcttctgt	ggctggcctg	atgatcacca	ccgaatgcac	ggttaccgac	2040
ctgccgaaaa	acgatgcagc	tgacttaggc	gctgctggcg	gtatgggagg	catgggtggc	2100
atgggaggca	tgatgtaatt	gccctgcacc	tcgcagaaat	aaacaaaccc	ccggg	2155

<210> 14

<211> 3139

<212> DNA

<213> Escherichia coli

<400> 14

atgggtaaaa	taattgggtat	cgacctgggt	actaccaact	cttgtgtagc	gattatggat	60
ggcaccactc	ctcgcgtgct	ggagaacgcc	gaaggcgatc	gcaccacgcc	ttctatcatt	120
gcctataccc	aggatggtga	aactctagtt	ggtcagccgg	ctaaacgtca	ggcagtgcag	180
aacccgcaaa	acactctgtt	tgcgattaaa	cgccgtgattg	gtcgccgctt	ccaggacgaa	240
gaagtacagc	gtgatgtttc	catcatgccg	ttcaaaatta	ttgctgctga	taacggcgac	300
gcatgggtcg	aagttaaagg	ccagaaaaatg	gcaccgcgcg	agatttctgc	tgaagtgtctg	360
aaaaaaatga	agaaaaacgc	tgaagattac	ctgggtgaac	cggttaactga	agctggtatc	420
accgtaccgg	catactttta	cgatgctcag	cgtcaggcaa	ccaaagacgc	aggccgtatc	480
gctggtctgg	aagtaaaacg	tatcatcaac	gaaccgaccg	cagctgcgct	ggcttacggt	540
ctggacaaa	gcactggcaa	ccgtactatc	gcggtttatg	acctgggtgg	tggtactttc	600
gatattttcta	ttatcgaaat	cgacgaagtt	gacggcgaaa	aaaccttcga	agttctggca	660
accaacgggtg	ataccacact	gggggggtgaa	gacttcgaca	gccgtctgat	caactatctg	720
gttgaagaat	tcaagaaaaga	tcagggcatt	gacctgcgca	acgatccgct	ggcaatgcag	780
cgctgaaaag	aaagcggcaga	aaaagcgaaa	atcgaaactgt	cttccgctca	gcagaccgac	840
gttaacctgc	catacatcac	tgcagacggg	accggtccga	aacacatgaa	catcaaagtg	900
actcgtgcga	aactggaaa	cctggttgaa	gatctggtaa	accgttccat	tgagccgctg	960
aaagttgcac	tgcaggacgc	tggcctgtcc	gtatctgata	tcgacgacgt	tatcctcggt	1020
gggtgtcaga	ctcgtatgcc	aatggttcag	aagaaagtgtg	ctgagttctt	tggtaaaagag	1080
ccgcgtaaaag	acgttaaccc	ggacgaagct	gtagcaatcg	gtgctgctgt	tcagggtggt	1140
gttctgactg	gtgacgtaaa	agacgtactg	ctgctggacg	ttaccccgct	gtctctgggt	1200
atcgaaacca	tggcggtgtg	gatgacgacg	ctgatcgoga	aaaacaccac	tatcccgacc	1260
aagcacagcc	aggtgttctc	taccgctgaa	gacaaccagt	ctgcggtaac	catccatgtg	1320
ctgcagggtg	aacgtaaaacg	tgcggctgat	aacaaatctc	tgggtcagtt	caacctagat	1380
ggtatcaacc	cggcaccgcg	cggcatgccg	cagatcgaa	ttaccttcga	tatcgatgct	1440
gacggtatcc	tgcacgtttc	cgcgaaagat	aaaaacagcg	gtaaagagca	gaagatcacc	1500
atcaaggctt	cttctggtct	gaacgaagat	gaaatccaga	aaatggtagc	cgacgcagaa	1560
gctaaccggg	aagctgaccg	taagtttgaa	gagctggtac	agactcgcaa	ccaggggcag	1620
catctgctgc	acagcaccgg	taagcaggtt	gaagaagcag	gcgacaaaact	gccggctgac	1680
gacaaaaactg	ctatcgagtc	tgcgctgact	gcactggaaa	ctgctctgaa	aggtgaagac	1740
aaagccgcta	tcgaagcgaa	aatgcaggaa	ctggcacagg	tttcccgaaa	actgatggaa	1800
atcgcccgac	agcaacatgc	ccagcagcag	actgccgggtg	ctgatgcttc	tgcaaacac	1860
gcgaaagatg	acgatgttgt	cgacgctgaa	tttgaagaag	tcaaaagacaa	aaaataatcg	1920
ccctataaac	gggtaattat	actgacacgg	gcgaagggga	atttcctctc	cgcggctgca	1980
ttcatctagg	ggcaatttaa	aaaagatggc	taagcaagat	tattacgaga	ttttaggcgt	2040
ttccaaaaca	gcggaagagc	gtgaaatcag	aaaggcctac	aaacgcctgg	ccatgaaata	2100
ccaccgggac	cgtaaccagg	gtgacaaa	ggccgaggcg	aaattttaa	agatcaagga	2160

```

agcttatgaa gttctgaccg actcgcaaaa acgtgcggca tacgatcagt atgggtcatgc 2220
tgcggtttgag caaggtggca tgggcggcgg cggttttggc ggcgggcgag acttcagcga 2280
tatttttggg gacgttttcg gcgatatatt tggcggcgga cgtggtcgtc aacgtgcggc 2340
gcgcgggtgct gatttacgct ataacatgga gctcaccctc gaagaagctg tacgtggcgt 2400
gaccaaagag atccgcattc cgactctgga agagtgtgac gtttgccacg gtagcgggtgc 2460
aaaaccagggt acacagccgc agacttgtcc gacctgtcat ggttctggtc aggtgcagat 2520
gcgccaggga ttcttcgctg tacagcagac ctgtccacac tgtcagggcc gcggtacgct 2580
gatcaaagat ccgtgcaaca aatgtcatgg tcatggctgt gttgagcgca gcaaaacgct 2640
gtccgttaaaa atcccggcag ggggtggacac tggagaccgc atccgtcttg cgggcgaagg 2700
tgaagcgggc gagcatggcg caccggcagg cgatctgtac gttcagggtc aggttaaaca 2760
gcacccgatt ttcgagcgtg aaggcaacaa cctgtattgc gaagtcccga tcaacttcgc 2820
tatggcggcg ctgggtggcg aaatcgaagt accgaccctt gatggctcgc tcaaactgaa 2880
agtgcctggc gaaacccaga ccggttaagc attccgtatg cgcggtaaaag gcgtcaagtc 2940
tgtccgcggg ggcgcacagg gtgatttgct gtgccgcgtt gtcgtcgaaa caccggtagg 3000
cctgaacgaa aggcagaaac agctgctgca agagctgcaa gaaagcttcg gtggcccaac 3060
cggcgagcac aacagcccgc gctcaaagag cttctttgat ggtgtgaaga agttttttga 3120
cgacctgacc cgctaataa 3139

```

<210> 15

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 15

cccccccggg atgggtaaaa taattggtat cgac

34

<210> 16

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 16

cgcgggatcc ttattagcgg gtcaggctgt caaaaaa

37

<210> 17

<211> 594

<212> DNA

<213> Escherichia coli

<400> 17

```

atgagtagta aagaacagaa aacgcctgag gggcaagccc cggaagaaat tatcatggat 60
cagcacgaag agattgagcg agttgagcca gaagcttctg ctgagcaggt ggatccgcgc 120
gatgaaaaag ttgcgaatct cgaagctcag ctggctgaag cccagaccgc tgaacgtgac 180
ggcattttgc gtgtaaaagc cgaatggaa aacctgcgtc gtcgtactga actggatatt 240
gaaaaagccc acaaattcgc gctggagaaa ttcatacaac aattgctgcc ggtgattgat 300
agcctggatc gtgcgctgga agtggtgat aaagctaacc cggatatgtc tgcgatgggt 360
gaaggcattg agctgacgct gaagtcgat ctggatggtg tgcgtaagtt tggcgttgaa 420
gtgatcgccg aaactaacgt cccactggac ccgaatgtgc atcaggccat cgcaatggtg 480
gaatctgatg acgttgccgc aggtaacgta ctgggcatta tgcagaaggg ttatacgtg 540
aatggtcgta cgattcgtgc ggcgatgggt actgtagcga aagcaaaagc ttaa 594

```

<210> 18

<211> 34

<212> DNA

<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 18  
cgcggaattc atgagtagta aagaacagaa aacg 34

<210> 19  
<211> 37  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 19  
aaaactgcag ttattaagct ttgtcttcg ctacagt 37

<210> 20  
<211> 2574  
<212> DNA  
<213> Escherichia coli

<400> 20  
atgctgtctg atcgtcttac taataaattc cagcttgctc ttgccgatgc ccaatcactt 60  
gcactcgggc acgacaacca atttatcgaa ccacttcatt taatgagcgc cctgctgaat 120  
cagggaagggg gtccggttag tccttttatta acatccgctg gcataaatgc tggccagttg 180  
cgcacagata tcaatcaggc attaaatcgt ttaccgcagg ttgaaggtag tgggtggtgat 240  
gtccagccat cacaggatct ggtgocggtt cttaatcttt gcgacaagct ggcgcaaaaa 300  
cgtggtgata actttatctc gtcagaactg ttctgtcttg cggcacttga gtctcgcggc 360  
acgctggccg acatcctgaa agcagcaggg gcgaccaccg ccaacattac tcaagcgatt 420  
gaacaaatgc gtggagggtga aagcgtgaac gatcaagggt ctgaagacca acgtcaggct 480  
ttgaaaaaat ataccatcga ccttacccga cgagccgaac agggcaaaact cgatccggtg 540  
attggtcgtg atgaagaaat tcgcgcgtacc attcaggtgc tgcaacgtcg tactaaaaat 600  
aaccgggtac tgattggtga acccgcgctc ggtaaaaact ccactcgttg aggtctggcg 660  
cagcgtatta tcaacggcga agtgccggaa ggggtgaaag gccgcgggt actggcgctg 720  
gatatgggag cgctggtggc tggggcgaaa tatcgcggtg agtttgaaga acgtttaaaa 780  
ggcgtgctta acgatcttgc caaacaggaa ggcaacgtca tcctatttat cgacgaatta 840  
cataccatgg tcggcgcggg taaagccgat ggcgcaatgg acgcccgaaa catgctgaaa 900  
ccggcgctgg cgcgtggtga attgcactgc gtaggtgcca cgacgcttga cgaatatcgc 960  
cagtacattg aaaaagatgc tgcgtggaa cgtcgtttcc agaaagtgtt tgttgccgag 1020  
ccttctgttg aagataccat tgcgattctg cgtggcctga aagaacgtta cgaattgcac 1080  
caccatgtgc aaattactga cccggcaatt gttgcagcgg cgacgttgct tcactcgctac 1140  
attgctgacc gtcagctgcc ggataaagcc atcgacctga tcgatgaagc agcatccagc 1200  
attcgtatgc agattgactc aaaaccagaa gaactcgacc gactcgatcg tcgtatcac 1260  
cagctcaaac tggaacaaca ggcgttaatg aaagagctcg atgaagccag taaaaaacgt 1320  
ctggatatgc tcaacgaaga actgagcgac aaagaacgtc agtactccga gttagaagaa 1380  
gagtggaaag cagagaaggc atcgctttct ggtacgcaga ccattaaagc ggaactggaa 1440  
caggcgaaaa tcgctattga acaggctcgc cgtgtggggg acctggcgcg gatgtctgaa 1500  
ctgcaatacg gcaaaatccc ggaactggaa aagcaactgg aagccgcaac gcagctcgaa 1560  
ggcaaaaacta tgcgtctgtt gcgtaataaa gtgaccgacg ccgaaattgc tgaagtgtctg 1620  
gcgcgttgga cggggattcc ggtttctcgc atgatggaaa gcgagcgcgaa aaaactgctg 1680  
cgtatggagc aagaactgca ccctgcgcta attggtcaga acgaagcggg tgatgcggta 1740  
tctaacgcta ttcgtcgtag ccgtgcgggg ctggcggtat caaatcgccc gattggttca 1800  
ttcctgttcc tcggcccaac tgggtgtggg aaaacagagc tttgtaaggc gctggcgaa 1860  
tttatgtttg atagcgacga ggcgatgggt gctatcgata tgtccgagtt tatggagaaa 1920  
cactcgtgtg ctcgtttggg ttgtgcgctt ccgggatatg tcggttatga agaaggtggc 1980  
tacctgacgc aagcgggtgc tcgtcgtccg tattccgtca tcctgctgga tgaagtggaa 2040  
aaagcgcac cggatgtctt caacattctg ttgcaggtac tggatgatgg gcgtctgact 2100  
gacgggcaag ggagaaacgg cgacttccgt aatacggctc tcattatgac ctctaacctc 2160  
ggttccgatc tgattcagga acgcttcggg gaactggatt atgcgcacat gaaagagctg 2220  
gtgctcgggt tggttaagca taacttccgt ccggaattca ttaaccgtat cgatgaagtg 2280  
gtggtcttcc atccgctggg tgaacagcac attgcctcga ttgcgcagat tcagttgaaa 2340  
cgtctgtaca aacgtctgga agaacgtggt tatgaaatcc acatttctga cgaggcgctg 2400

aaactgctga gcgagaaagg ttaogatccg gtctatggg cagtcctct gaaacgtgca 2460  
attcagcagc agatcgaaaa cccgctggca cagcaaatac tgtctgggta attgggtccg 2520  
ggtaaagtga ttgcctgga agttaatgaa gaccggattg tcgccgtcca gtaa 2574

<210> 21  
<211> 34  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 21  
aaaactgcag atgcgtctgg atcgtcttac taat 34

<210> 22  
<211> 37  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 22  
cccgggaagc ttattactgg acggcgacaa tccgggc 37

bioRxiv preprint doi: <https://doi.org/10.1101/000000>; this version posted January 1, 2014. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.